



**South  
Cambridgeshire  
District Council**



## Short term air quality in Harston

---

August 2021



## Executive Summary

Air quality was monitored in Harston using new Zephyr monitoring technology in the period January – June 2021. Harston was selected as a pilot monitoring location due to local concerns about traffic and the nearby travel hub development, and monitoring was carried out outside Harston and Newton Community Primary School, as it is recognised that children are among the most vulnerable to the impacts of air pollution. It was found that concentrations of the main pollutants of concern, nitrogen dioxide (NO<sub>2</sub>) and particulate matter (PM<sub>10</sub> and PM<sub>2.5</sub>), were comfortably below the national objectives for annual mean concentrations and there were no exceedances of the short-term objectives, representing good air quality. This is in line with long-term concentrations measured across the South Cambridgeshire district and reflects the rural nature of the area. South Cambridgeshire residents can help to improve local air quality through actions such as reducing idling of car engines and increasing walking and cycling where possible. This report can be read alongside the yearly Air Quality Annual Status Report (ASR) and future reports of other localised studies, which will be available on our [website](#).



## Glossary

**Annualisation** – a calculation process used to estimate an average concentration for a full year from a shorter period.

**Annual mean** – the average concentration across a full calendar year.

**AQMA – Air Quality Management Area** – an area where air pollutant concentrations exceed or are likely to exceed the relevant air quality objectives. AQMAs are declared for specific pollutants and objectives.

**Continuous monitor/monitoring station** – instruments which measure air pollution all the time and therefore can give a concentration attributed to a specific time.

**Diffusion tube** – Small plastic tube containing a metal mesh which is coated with a chemical that absorbs nitrogen dioxide. This is exposed to the air in a fixed location for a known amount of time, usually a month, and then sent to a lab for analysis. This provides an average concentration for the time it is exposed.

**Nitrogen dioxide (NO<sub>2</sub>)** – a gas predominantly formed following the burning of fossil fuels, which can cause irritation of the airways and exacerbate symptoms of other conditions.

**Particulate matter (PM<sub>2.5</sub> and PM<sub>10</sub>)** – the number refers to the size of the particulates in micrometres (one millionth of a metre) – a mix of solid particles and liquid droplets of various sizes and composition, the smallest of which can get into the blood and be transported around the body.

**Real-time monitoring** – see also continuous monitoring – monitoring which takes place at regular intervals all the time and therefore can give a concentration attributed to a specific time.

**µg/m<sup>3</sup>** – micrograms per cubic metre, the standard units of measurement of air pollutants including nitrogen dioxide and particulate matter.

**Zephyr** – a type of relatively compact and lightweight air pollution sensors that measure harmful gases and particle matter in real-time.



# Update on pilot Zephyr monitor in Harston

## Introduction

### Purpose of this report


This is a report to provide an update on the short-term air quality monitoring pilot study in Harston using new Zephyr monitoring technology. Monitoring was carried out in the period January – June 2021. The study was designed to be a short-term study monitoring air quality outside Harston and Newton Community Primary School following concerns about traffic and the nearby travel hub development, whilst also acting as a pilot scheme to test new air quality monitoring equipment. It also serves to create additional local awareness of air quality in our area and enable people to make informed choices around how they can impact on improving air quality in their area.

### Air Quality in South Cambridgeshire

South Cambridgeshire is a rural district which enjoys generally good air quality, with both short-term and long-term pollution levels below the national objectives at all monitored locations. This means we benefit from cleaner air to breathe and less pollution related health problems. The area is undergoing significant growth with major developments to keep up with the increase in demand for housing, including Northstowe (10,000 dwellings), Waterbeach Barracks (6000-10,000 dwellings), Bourn Airfield and Cambourne West, shown in Figure 1. Air quality impacts in the district are mainly related to these areas of growth and the major roads running through the district, including the A14 and M11/A11 corridors, and therefore this remains an important issue.



Figure 1 – Location of Major Development sites in South Cambridgeshire



Air quality is an important topic as air pollution can impact our health, particularly affecting the most vulnerable, including children and those with underlying conditions. Air quality is monitored across the district using a network of diffusion tubes and continuous monitoring stations, which provide accurate air quality measurements in real-time, in addition to the new Zephyr monitors to be used for short term monitoring. For more information and detail on the importance of air quality and air quality in South Cambridgeshire, please refer to Appendix 1 – Air Quality Frequently Asked Questions or visit our [website](#). Additionally, ideas on how anyone can play a role in improving local air quality can be found in Appendix 2 – How to get Involved with Local Air Quality.

### The ‘Zephyr’ Air Quality Sensor

Zephyr monitors are compact and lightweight air pollution sensors that measure harmful gases and particle matter in real-time, including the main pollutants of



concern (NO<sub>2</sub> and PM<sub>10</sub> and PM<sub>2.5</sub> particulate matter). They can run off internal batteries or be powered by a solar panel and can therefore be fixed in a specific location, mostly commonly a lamp post, or used as a mobile monitor. The sensors provide detailed air quality measurements in real-time to help identify pollution hotspots at a localised level, for example busy junctions. Other potential studies include investigating air quality around school and looking into the impacts of wood burning stoves. Zephyr sensors can be used in isolation individually or deployed as a network of sensors across a wider area to build up a more detailed picture.<sup>1</sup>

The data from a Zephyr sensor cannot be treated with the same confidence as that from one of our continuous monitor stations, where the data is 'ratified' after checks, however it has been shown to provide accurate indicative measurements and is therefore appropriate for a wide range of studies, including this.

### Monitoring Location

Harston was selected as a pilot location for the Zephyr sensor due to local ongoing development and infrastructure projects, such as the nearby travel hub site. The instrument is located near to Harston & Newton Community Primary School to additionally contribute to a council study looking into air quality around schools.

The monitor is located on the High Street of Harston, close to Harston and Newton Community Primary School and is measuring the main pollutants of concern, nitrogen dioxide (NO<sub>2</sub>) and particulate matter (PM<sub>10</sub> and PM<sub>2.5</sub>), among others. The location of the Zephyr can be seen on Figure 2, below, as well as the location of the diffusion tube located in Harston (DT8N). One of the main reasons for selecting a primary school as the location is that children are amongst the most vulnerable to the effects of air pollution, which was reflected by the theme of [Clean Air Day](#) in 2021 of 'protect our children's health from air pollution'<sup>2</sup>. This is due to children's airways and respiratory systems being less developed than an adult's and because they breathe more rapidly than adults.


---

<sup>1</sup> <https://www.earthsense.co.uk/zephyr>

<sup>2</sup> <https://www.cleanairday.org.uk/>



Figure 2 - Location of the pilot Zephyr sensor in Harston





## Monitoring Data and Comparison with Objectives

The average monthly concentrations measured in the period January to June 2021 are shown in Table 1, below, with the annual mean objective shown for information. This data is also represented in Figure 3. Data was also recorded for the month of December, however due to an issue with the instrument power for a significant period of this month there was an extended 'settling in' period, meaning large sections of the NO<sub>2</sub> data were invalid. This was confirmed as invalid by comparison to the particulate matter measurements, data from the diffusion tubes in Harston in December and the data from other continuous monitor locations in the district in the same time period. Consequently, it was decided not to include the December data in the report, as per the Defra guidance in Technical Guidance TG16<sup>3</sup>, and extend the study until the end of June so a full 6 months of data would be recorded and available.


Table 1 – Zephyr Air Quality data – monthly average concentrations

Month	Pollutant monthly average concentration / $\mu\text{g}/\text{m}^3$		
	NO <sub>2</sub>	PM <sub>10</sub>	PM <sub>2.5</sub>
January 2021	5.7	14.5	7.3
February 2021	10.3	14.4	8.3
March 2021	15.1	17.0	10.1
April 2021	20.7	13.0	9.5
May 2021	16.3	10.6	7.5
June 2021	12.0	11.7	8.7
Objective (annual mean)	40	40	N/A

<sup>3</sup> Defra Local Air Quality Management (LAQM) Technical Guidance TG(16), 2018, <https://laqm.defra.gov.uk/documents/LAQM-TG16-February-18-v1.pdf>



Figure 3 – Zephyr Air Quality data – monthly average concentrations and national annual mean objective



With the exception of NO<sub>2</sub> data around January, which was lower than seen elsewhere, the data and trends recorded by the Harston Zephyr are generally consistent with those seen at other monitoring locations across the district. This suggests that overall there can be confidence in the data collected, although there may be a longer ‘settling in’ period for the NO<sub>2</sub> sensor. Due to this it could be worth considering monitoring at the same location in the future.

In addition, the average concentrations of each pollutant for the whole six-month period January – June were calculated and then ‘annualised’ to give estimated annual mean concentrations to allow better comparison to the annual mean objectives. Annualisation is a calculation process used to estimate an average concentration for a full year from a shorter period, such as the 6 months in this study. This is done to avoid the annual average being influenced by short-term events or seasonal changes, such as one day of high pollution like bonfire night or pollution concentrations often being higher in the winter than the summer. The data was annualised using 2020 data from a range of continuous monitoring background site and is shown in Table 2, below. Full annualisation details are available in Appendix 3 – Annualisation of short-term data. The annual mean NO<sub>2</sub> concentration measured at




the diffusion tube (DT8N) located in Harston for 2020 was also included for comparison.

Table 2 – Zephyr Air Quality Data – annualised annual mean concentrations

	Pollutant monthly average concentration / $\mu\text{g}/\text{m}^3$		
	NO <sub>2</sub>	PM <sub>10</sub>	PM <sub>2.5</sub>
Measured data average Jan-June	13.4	13.6	8.6
Annualisation factor	0.97	0.96	0.94
<b>Annualised annual mean – Harston Zephyr</b>	<b>13.0</b>	<b>13.0</b>	<b>8.1</b>
DT8N (2020)	12.3	N/A	N/A
Objective (annual mean)	40	40	N/A

As shown in Table 1 and Table 2, the long-term annual mean concentrations of the main pollutants of concern at the Harston Zephyr are significantly below the national objectives. For NO<sub>2</sub> this is supported by the data from the diffusion tube in 2020. The NO<sub>2</sub> data is shown in also displayed on Figure 4, below.

Figure 4 – NO<sub>2</sub> annual mean concentrations after annualisation and annual mean objective





The Zephyr also allows measurements of the short-term concentrations of pollutants, which are studied through 1-hour means for NO<sub>2</sub> and 24-hour means for PM<sub>10</sub>.

These are presented and compared to the national objectives in Table 3, below. The short-term objectives are presented as hourly/daily concentrations that should not be exceeded more than a certain number of times in a year. There is currently no short-term objective for PM<sub>2.5</sub>.

Table 3 – Zephyr Air Quality data – short-term average concentrations

Month	Number of exceedances of short-term objective	
	NO <sub>2</sub> 1-hour mean	PM <sub>10</sub> 24-hour mean
January 2021	0	0
February 2021	0	0
March 2021	0	0
April 2021	0	0
May 2021	0	0
June 2021	0	0
Objective	200 µg/m <sup>3</sup> *	50 µg/m <sup>3</sup> **

\*Not to be exceeded more than 18 times a year

\*\*Not to be exceeded more than 35 times per year

As shown in Table 3, there were no exceedances of the short-term objectives for NO<sub>2</sub> or PM<sub>10</sub>. The relevant maximum short-term concentrations of the pollutants were also recorded. For NO<sub>2</sub> the maximum 1-hour concentration measured during the six-month period was 76.6 µg/m<sup>3</sup>, which occurred during the evening rush hour on a weekday in March; this is well under the 200 µg/m<sup>3</sup> threshold. For PM<sub>10</sub>, the maximum 24-hour concentration recorded was 49.1 µg/m<sup>3</sup>, below the 50 µg/m<sup>3</sup> objective which should not be exceeded more than 35 times. This was the only 24-



hour average above  $45 \mu\text{g}/\text{m}^3$ , with only one additional 24-hour period above  $40 \mu\text{g}/\text{m}^3$ , which occurred the day before.

## Summary

The data measured by the Zephyr real-time monitor in the period January – June 2021 shows that the air quality in Harston remains good, with estimated annual mean concentrations (as well as the measured monthly averages) of all the main pollutants of concern well below the national objectives, and no exceedances of the short-term national objectives. In general, the data from the Zephyr was consistent with that from the rest of the monitoring sites in the district during the monitoring period, which provides confidence in the instrument and data collected. This also matches the general patterns seen across the South Cambridgeshire district of good air quality. However, due to the importance of air quality and its links to health, it remains important to both monitor air quality across the district and take actions to improve air quality in our area. Ideas on how to play a role in improving local air quality can be found in Appendix 2 – How to get Involved with Local Air Quality.



## Appendix 1 – Air Quality Frequently Asked Questions

### Why is air quality important?

There are a number of reasons air quality is important. In particular, polluted air is the biggest environmental threat to health in the UK. It is linked to up to 36,000 deaths per year from long-term exposure<sup>4</sup>. The main impacts of poor air quality are contributing to heart and lung conditions, but air quality has also been linked to a wide range of issues<sup>5</sup>. Air pollution also particularly effects the most vulnerable, including children and older people and those with existing lung and heart conditions. Air quality also strongly links to climate change, as many of the causes of the issues are the same, such as the burning of fossil fuels. This means that actions taken to improve air quality also helps prevent climate change.

### How does the Council monitor air quality?

South Cambridgeshire District Council operates a monitoring network of over 30 locations across the district, made up of diffusion tubes and three continuous monitoring sites, which measure air quality accurately in real-time. This existing monitoring network allows the long-term monitoring of trends and changes in air quality across the district. Live data from the three continuous monitoring stations are available at <https://scambs-airquality.ricardo-aea.com/>. In addition, the Council has purchased three Zephyr air quality sensors which provide real-time measurements for the main pollutants of concern from a single monitor. These can be used for shorter-term monitoring to identify hotspots of pollution or be used in a range of targeted studies to complement our existing monitoring network. The first of these instruments was installed in Harston, with subsequent monitors installed in Cambourne and Northstowe.

---

<sup>4</sup> Defra. Air quality appraisal: damage cost guidance, July 2020

<sup>5</sup> Public Health England. Air Quality: A Briefing for Directors of Public Health, 2017



## What else does the Council do around air quality?

As well as monitoring air quality, the Council acts to improve air quality through its Green to the Core focus, including an air quality strategy designed to go beyond simply meeting the national objectives, Zero Carbon Community Grants to fund community initiatives to improve sustainability, such as encouraging and enabling cycling which in turn helps air quality, and by considering air quality during the planning process<sup>6,7</sup>. Ideas on how anyone can play a role in improving local air quality can be found in Appendix 2 – How to get Involved with Local Air Quality.

## What are the main pollutants of concern?

The main pollutants of concern are:

- Nitrogen Dioxide (NO<sub>2</sub>) – a gas predominantly formed following the burning of fossil fuels, which can cause irritation of the airways and exacerbate symptoms of other conditions
- Particulate Matter (PM<sub>10</sub> and PM<sub>2.5</sub>), where the number refers to the size of the particulates in micrometres – a mix of solid particles and liquid droplets of various sizes and composition, the smallest of which can get into the blood and be transported around the body<sup>8</sup>

## What are the air quality objectives?

For these pollutants national objective levels have been set which must be achieved by local authorities, otherwise an Air Quality Management Area (AQMA) must be declared for the objective which is being exceeded. Objectives have been set for both long-term concentrations (measured as annual means) and short-term concentrations (hourly means for NO<sub>2</sub> and daily means for PM<sub>10</sub>). South Cambridgeshire District Council currently has one AQMA, along the A14 between Bar Hill and Milton, which was declared in 2008 for NO<sub>2</sub> annual mean and PM<sub>10</sub> 24-hour mean. It is proposed to revoke this AQMA in 2021 due to sustained compliance

---

<sup>6</sup> Being green to our core <https://www.scambs.gov.uk/your-council-and-democracy/performance-and-plans/our-business-plan/>

<sup>7</sup> Zero Carbon Communities Grant <https://www.scambs.gov.uk/community-development/grants/zero-carbon-communities-grant/>.

<sup>8</sup> Defra, Clean Air Strategy, 2019



with the relevant objectives in line with Defra guidance and the Council's constitution. The Air Quality Objectives for England are set out in Table 4. In addition, local authorities are expected to work towards reducing emissions and concentrations of PM<sub>2.5</sub> (particulate matter with a diameter of 2.5 µm or less), although there is currently no legal objective.

Table 4 – Air Quality Objectives in England

Pollutant	Air Quality Objective – Concentration	Air Quality Objective – Measured as
Nitrogen Dioxide (NO <sub>2</sub> )	200 µg/m <sup>3</sup> not to be exceeded more than 18 times a year	1-hour mean
Nitrogen Dioxide (NO <sub>2</sub> )	40 µg/m <sup>3</sup>	Annual mean
Particulate Matter (PM <sub>10</sub> )	50 µg/m <sup>3</sup> , not to be exceeded more than 35 times a year	24-hour mean
Particulate Matter (PM <sub>10</sub> )	40 µg/m <sup>3</sup>	Annual mean
Sulphur Dioxide (SO <sub>2</sub> )	350 µg/m <sup>3</sup> , not to be exceeded more than 24 times a year	1-hour mean
Sulphur Dioxide (SO <sub>2</sub> )	125 µg/m <sup>3</sup> , not to be exceeded more than 3 times a year	24-hour mean
Sulphur Dioxide (SO <sub>2</sub> )	266 µg/m <sup>3</sup> , not to be exceeded more than 35 times a year	15-minute mean

### If air pollution is a result of vehicles utilising the A14, how can local residents change this?

There are a number of way local residents can have an impact on air quality through everyday actions, such as those mentioned in **Error! Not a valid bookmark self-reference..** Many of these are very small changes that can add up to a big impact.



## Appendix 2 – How to get Involved with Local Air Quality

Annual reports and details on air quality monitoring are available on our website, <https://www.scambs.gov.uk/environment/pollution/air-pollution/local-air-quality-management/>, and you can share your views via our email address, [air.quality@scambs.gov.uk](mailto:air.quality@scambs.gov.uk).

Although air quality in the South Cambridgeshire District is generally good, with concentrations below the objectives, there are actions we can all take to improve it further. Ways you can help to improve air quality in South Cambs include:

- Minimise car use wherever possible:
  - Avoid using your car for short trips (under 2 miles) - short trips are very polluting as modern engines need to reach a very high temperature to work efficiently; on short trips it won't reach that temperature.
  - For short journeys try cycling or walking more often – this helps you stay healthy and saves you money in fuel costs.
  - For longer journeys consider public transport options.
  - Use journey-planning apps such as MyBusTrip or MotionMap for travel by bus, train, walking and cycling.
- Switch it off – don't leave your car engine idling if you are stationary e.g. waiting to pick someone up, in a traffic jam or waiting at level crossings.
- When driving, use techniques that help you use less fuel, like driving more slowly and smoothly.
  - You could use 10% less fuel by following the tips on the AA website [http://www.theaa.com/motoring\\_advice/fuels-and-environment/drive-smart.html](http://www.theaa.com/motoring_advice/fuels-and-environment/drive-smart.html).
  - Like switching your engine off when stationary, this will not only reduce your emissions of air pollution but will save fuel and therefore money too!
- Consider making your next vehicle an electric vehicle.
- Join a car club or car-share regularly.
- Consider working at home where possible – the first Covid-19 lockdown showed widespread improvements in the air quality as the amount of people travelled reduced.



- Use less energy at home – consider a smart meter to monitor usage and be aware of boiler standards.
- Opt for ‘green energy’ tariffs where available or switch to renewable sources of heating or power.
- Reduce the use of solid fuel stoves and open fires – domestic burning is now the single biggest source of particulate matter pollution in the UK (greater than traffic and industry).
  - If you are burning wood or coal ensure any fuel used meets the new standards of moisture content and emissions – more information is available at <https://woodsurre.co.uk/are-you-ready-to-burn/>
- Make your children aware of the impact that day to day activities have on air quality.



## Appendix 3 – Annualisation of short-term data

Annualisation is a calculation process used to estimate an average concentration for a full year from a shorter period, such as the 6 months in this study. Annualisation ratios are worked out as a ratio of the average concentration in a full year (annual mean (Am)) to the average in the actual monitoring period measured (period mean (Pm)), using data from background continuous sites. The average concentration from the Zephyr data during the monitoring period is then multiplied by that ratio to give an estimate of the average concentration at the Zephyr for a full year.

The data from the period January to June 2021 was annualised according to the process set out in box 7.9 of Defra's Local Air Quality Management Technical Guidance (TG16). Continuous monitoring background sites were used for the annualisation calculations. Full year data for 2021 is not yet available therefore 2020 data was used for the annual mean concentrations.

NO<sub>2</sub>:

Background Site	Annual mean (Am)	Period mean (Pm)	Ratio (Am/Pm)
Orchard Park	10.6	11.4	0.93
Wicken Fen	6.7	6.2	1.08
Northampton Spring Park	9.3	10.6	0.88
Norwich Lakenfields	9.8	9.9	0.99
Average ratio	-	-	0.97

PM<sub>10</sub>:

Background Site	Annual mean (Am)	Period mean (Pm)	Ratio (Am/Pm)
Orchard Park	12.2	12.3	0.99
Norwich Lakenfields	12.8	13.9	0.92
Average ratio	-	-	0.96



PM<sub>2.5</sub>:

Background Site	Annual mean (Am)	Period mean (Pm)	Ratio (Am/Pm)
Orchard Park	12.5	12.8	0.98
Northampton Spring Park	10.3	11.0	0.94
Norwich Lakenfields	8.3	9.2	0.90
Average ratio	-	-	0.94